

The 2-D electron gas at arbitrary spin polarizations and arbitrary coupling strengths: exchange-correlation energies, distribution functions and spin-polarized phases.

François Perrot, M.W.C. Dharma-wardana[†]

National Research Council, Ottawa, Canada. K1A 0R6

Centre d'Etudes de Bruyères le Châtel, P. O. Box 12, 91689 Bruyères le Châtel, France

(24 May 1999)

We use a recent approach [Phys. Rev. Letters, **84**, 959 (2000)] for including Coulomb interactions in quantum systems via a classical mapping of the pair-distribution functions (PDFs) for a study of the 2-D electron gas. As in the 3-D case, the “quantum temperature” T_q of a *classical* 2-D Coulomb fluid which has the same correlation energy as the quantum fluid is determined as a function of the density parameter r_s . Spin-dependent exchange-correlation energies are reported. Comparisons of the spin-dependent pair-distributions and other calculated properties with any available 2-D quantum Monte Carlo (QMC) results show excellent agreement, strongly favouring more recent QMC data. The interesting novel physics brought to light by this study are: (a) the independently determined quantum-temperatures for 3-D and 2-D are found to be approximately the *same*, (i.e., universal) function of the classical coupling constant Γ ; (b) the coupling Γ increases rapidly with r_s in 2-D, making it comparatively more coupled than in 3-D; the stronger coupling in 2-D requires bridge corrections to the hyper-netted-chain method which is adequate in 3-D; (c) the Helmholtz free energy of spin-polarized and unpolarized phases have been calculated. The existence of a spin-polarized 2-D liquid near $r_s \sim 30$, is found to be a marginal possibility. These results pertain to clean uniform 2-D electron systems.

PACS Numbers: 05.30.Fk, 71.10.+x, 71.45.Gm

The past three decades have established the two-dimensional electron gas (2DEG) as a goldmine of new physics and novel technologies.¹ The 2DEG, ranging from a classical to a quantum system, occurs on liquid-He surfaces, semiconductor hetero-interfaces, and in the CuO layers of high- T_c superconductors. The interest in the quantum Hall effect, the metal-insulator transition, Kosterlitz-Thouless type transitions,² and applications to nanotechnology and spintronics widen its importance. The exchange-correlation energy (E_{xc}) of the 2DEG is important in its own right as an input to the density-functional theory (DFT) of inhomogeneous 2-D systems.

The physics of the 2DEG depends crucially on the “coupling parameter” $\Gamma = (\text{potential energy})/(\text{kinetic energy})$ arising from the Coulomb interactions. The Γ for the 2DEG at $T = 0$ and mean density n is equal to the mean-disk radius $r_s = (\pi n)^{-1/2}$ per electron. In the absence of a magnetic field or disorder effects, the density parameter r_s , the spin polarization ζ and the temperature T are the only relevant physical variables. A major problem in the 2DEG is the need for a theoretical method applicable at *arbitrary* values of Γ , ζ and T . As in the 3-D electron gas (3DEG), various perturbation methods, equations-of-motion methods, e.g., those of Singwi et al (STLS),³ work for relatively weak coupling, but lead to negative pair-distribution functions (PDFs) even at moderate r_s . In spite of a large effort, the calculation of the 2DEG $g_{ij}(r)$, where i, j are spin species, at arbitrary spin-polarization ζ , r_s , and T is unsolved, except by direct quantum simulation techniques. Such “quantum Monte Carlo” (QMC) methods,⁴ as well as Feenberg

techniques,⁵ assume a trial wavefunction $\psi = FD$ where F is a correlation factor and D is a Slater determinant. The variational Monte Carlo (VMC) method,⁶ and the fixed-node Green-function Monte-Carlo method (GFMC) have been applied to the 2DEG, providing the exchange-correlation energy, $E_{xc}(r_s)$ at $T = 0$ in a parametrized form.⁶ Finite- T systems involve excited states as well as the ground state. While it is easy to get good $E_{xc}(r_s)$ at $T = 0$, $\zeta = 0$, the same is not true for most properties like the PDFs, local-field corrections (LFCs), and xc-energies at finite T , spin polarization ζ and at strong coupling.

Recently we presented a computationally simple, conceptually novel *classical* method for calculating the PDFs and other properties (e.g., static response) of the 3DEG at arbitrary coupling, spin polarization and temperature.⁹ The method is based on identifying a “quantum temperature” T_q such that the correlation energy of a corresponding *classical* Coulomb fluid at T_q is equal to that of the quantum fluid at $T = 0$. Our only “many-body” input is a set of values of $E_c(r_s)$ for determining T_q . Once $T_q(r_s)$ is known, many properties inaccessible to standard methods become readily calculable. A discussion of the $T = 0$ and finite- T 3DEG was presented earlier.^{9,10}

A method which works in 3-D does not necessarily work in reduced dimensions. A first objective of this letter is to show that the same ideas apply to the challenging problems of the 2DEG. In addition, this study reveals previously unexpected aspects of the quantum temperature T_q , exposes the long-range nature of the Pauli exclusion effect in the 2-D system, and provides new results for the partially polarized 2-D electron liquid. It is found

to be numerically precise enough, even at $r_s = 30$, to selectively favour newer QMC calculations.⁷ We also examine the stability of spin-polarized phases of the 2DEG at finite- T and sufficiently large r_s .

The PDFs of the 3DEG could be accurately calculated using the hyper-netted-chain (HNC) approximation.¹¹ The 2-D system requires that bridge-type cluster corrections to the HNC approximation be included. A comparison of the 2DEG and 3DEG forms of $T_q(r_s)$ reveals that the quantum temperature is approximately a universal function of the classical-fluid coupling constant.

The essence of our method is to start from the quantum mechanical $g^0(r)$ of the non-interacting problem and build up the interacting $g(r)$ by classical methods.

Consider a fluid of mean density n containing two spin species with concentrations $x_i = n_i/n$. We deal with the physical temperature T of the 2DEG, while the temperature T_{cf} of the classical fluid is $1/\beta$. Since the leading dependence of the energy on temperature is quadratic, we assume that $T_{cf} = \sqrt{(T^2 + T_q^2)}$. This is clearly valid for $T = 0$ and for high T , and was justified in more detail in ref. 10. In this letter the main effort is to study the 2DEG near $T = 0$ and at zero magnetic field by determining T_q . The equations for the PDFs of a classical fluid, and the Ornstein-Zernike(OZ) relations are:

$$g_{ij}(r) = \exp[-\beta\phi_{ij}(r) + h_{ij}(r) - c_{ij}(r) + B_{ij}(r)] \quad (1)$$

$$h_{ij}(r) = c_{ij}(r) + \sum_s n_s \int d\mathbf{r}' h_{i,s}(|\mathbf{r} - \mathbf{r}'|) c_{s,j}(\mathbf{r}') \quad (2)$$

Here $\phi_{ij}(r)$ is the pair potential between the species i, j . For two electrons this is just the Coulomb potential $V_{cou}(r)$. If the spins are parallel, the Pauli principle prevents occupation of the same spatial orbital. As before,⁹ we introduce a “Pauli potential”, $\mathcal{P}(r)$. Thus $\phi_{ij}(r)$ becomes $\mathcal{P}(r)\delta_{ij} + V_{cou}(r)$. The function $h(r) = g(r) - 1$ is related to the structure factor $S(k)$ by a Fourier transform. The $c(r)$ is the “direct correlation function (DCF)” of the OZ equations. The $B_{ij}(r)$ in Eq. 1 is the “bridge” term due to certain cluster interactions. If this is neglected, Eqs. 1-2 form a closed set defining the HNC approximation. The HNC is sufficient for the 3DEG for the range of r_s studied previously.⁹ The classical coupling constant of the 2DEG is found to increase with r_s more rapidly than in the 3DEG. Thus, although $\mathcal{P}(r)$ restricts clustering effects in the case of g_{ii} , bridge contributions (which contain irreducible 3-body and higher terms) are found to be important for g_{12} . The inclusion of a bridge term into the HNC equations, usually via a “hard-sphere” model,¹² is well known in 3-D problems, but is less unexplored for 2-D systems. In this context, it is noteworthy that Kwon et al. had to include three-body correlations in QMC trial functions.^{7,8}

In the non-interacting system at temperature T , $n = n_1 + n_2$, $x_i = n_i/n$, the anti- $\parallel$ $g_{12}^0(r, T)$ is unity while

$$h_{11}^0(\mathbf{r}) = -\frac{1}{n_i^2} \sum_{\mathbf{k}_1, \mathbf{k}_2} n(k_1) n(k_2) e^{i(\mathbf{k}_1 - \mathbf{k}_2) \cdot \mathbf{r}} = -[f(r)]^2$$

Here $\mathbf{k}, \mathbf{r}$ are 2-D vectors and $n(k)$ is the Fermi occupation number at the temperature T . At $T = 0$ $f(r) = 2J_1(k_i r)/k_r$ where $J_1(x)$ is a Bessel function. The Pauli exclusion potential is defined by:

$$\beta\mathcal{P}(r) = h_{11}^0(r) - c_{11}^0(r) - \ln[g_{11}^0(r)] \quad (3)$$

where, e.g., $c_{ii}^0(r)$ is the spin- $\parallel$ DCF of the O-Z equation. At $T = 0$ it may be shown that $\beta\mathcal{P}(r) \sim -2\ln(r)$ for $r \rightarrow 0$, $\beta\mathcal{P}(r) \sim \pi/rk_F$ for $r \rightarrow \infty$, $\beta\mathcal{P}(k) \sim -4\pi/k^2$ for $k \rightarrow \infty$. Only the product $\beta\mathcal{P}(r)$ is determined. The classical fluid “temperature” $T_{cf}=1/\beta$ is still undefined. The Pauli potential is a universal function of rk_F , and limited to about a thermal wavelength λ_{th} at finite T .

The next step in the method is to use the full $\phi_{ij}(r)$, and solve the coupled HNC and OZ equations for the binary (up and down spins) *interacting* fluid. The Coulomb potential $V_{cou}(r)$ for two point-charge electrons is $1/r$. However, an electron at the temperature T is localized to within a thermal wavelength. Thus, for the 3DEG we used a “diffraction corrected” form:¹³

$$V_{cou}(r) = (1/r)[1 - e^{-rk_{th}}] \quad (4)$$

where k_{th} was taken to be $k_{th}^0 = (2\pi m^* T_{cf})^{1/2}$ as in Minno et al.¹³ Here $m^*=1/2$ is the reduced mass of the electron *pair*. In the case of the 2DEG we have determined k_{th} by numerically solving the Schrodinger equation for a pair of 2-D electrons in the potential $1/r$ and calculating the electron density in each normalized state.¹⁴ Assuming that the diffraction correction has the same form as before, i.e., $V_d(r)=\exp(-k_{th}r)/r$, it was found that $k_{th}/k_{th}^0 = 1.158T^{0.103}$, where T is in au.

As in the 3DEG, we need the “quantum temperature” T_q for each r_s of the 2DEG, such that the classical 2D-fluid has the same $E_c(r_s)$ as the quantum fluid. Unlike in the 3DEG, an additional complication in the 2DEG is the need for a bridge function. In the $\parallel$ -spin case, if the hard-disk radius of the bridge function is smaller than the range of the Pauli potential, we may neglect the bridge term. In the anti- $\parallel$ case, we need a bridge term B_{12} explicitly included, say, via a hard-disk model.¹⁵ The same bridge term enters the paramagnetic PDF, $g(r) = 0.5(g_{11} + g_{12})$ via the g_{12} . We can decouple the determination of T_q and B_{12} by obtaining T_q from the *fully spin polarized* system as the bridge function is negligible in this case. Using the analytic fit of $E_c(r_s, \zeta = 1)$ given by Tanatar and Ceperley,⁶ $t = T_q/T_F$ is found to be fitted by:

$$t = 2/(1 + 0.86413(r_s^{1/6} - 1)^2) \quad (5)$$

This looks very different from the $T_q(r_s)$ mapping for the 3DEG. However, when T_q is plotted as a function of the *classical* coupling constant $\Gamma_c = 1/(T_q r_s)$, (cf. Table I), the 3DEG and 2DEG maps are seen to be almost a single universal function (Fig. 1.(a)). This result is *conceptually very interesting* since T_q is really a single-parameter representation of the density-functional correlation energy of the electron gas, irrespective of dimensionality.

Unlike in the 3DEG, we need the bridge function for g_{12} in order to implement the method. We include only the anti- $\parallel$ $B_{12}(r)$, and hence a one-component hard-disk packing-fraction η is needed. The effective coupling constant, $\Gamma_{eff} = \phi(r_s)/T_{cf}$, is a function of $r_s T_q$ and $k_{th} r_s$. That is, r_s/t and t^3/r_s where $t = T_q/T_F$. The Gibbs-Bogoliubov expression for the free energy of the one-component hard-disk system has the same dependence. Also, at large r_s the packing fraction should tend to a finite limit. Since t decreases as $r_s^{-1/3}$, we consider the simple functional form $\eta = \theta t r_s^{1/3}$. Here θ is a “global” parameter valid for the whole range of r_s and hence the procedure does not involve fitting at each r_s . A constant value, $\theta = 0.1175$, proved to be applicable to the range $r_s = 1 - 30$ studied here. Values of the packing fraction η , the QMC and classical-map HNC (CHNC) values of $E_c(r_s, \zeta = 0)$ for the unpolarized 2DEG are given in Table I. Our $E_c(r_s, \zeta = 1)$ agrees with Tanatar-Ceperley spin-polarized QMC by construction. It is common for η to approach ~ 0.49 at the liquid-solid transition. The trend in the value of η seen in Table I is consistent with the finding of Tanatar and Ceperley that the 2DEG becomes a Wigner solid for $r_s = 37 \pm 5$.

Using the T_q and the hard-disk bridge parameter θ , we can calculate $E_c(r_s, \zeta, T)$ at any ζ . Our results (at $T = 0$) can be presented via a “polarization function”:

$$p(r_s, \zeta) = \frac{E_c(r_s, \zeta) - E_c(r_s, 0)}{E_c(r_s, 1) - E_c(r_s, 0)} = \frac{\zeta_+^{\alpha(r_s)} - \zeta_-^{\alpha(r_s)} - 2}{2^{\alpha(r_s)} - 2}$$

$$\alpha(r_s) = C_1 - C_2/r_s + C_3/r_s^{2/3} - C_4/r_s^{1/3} \quad (6)$$

Here $\zeta_{\pm} = (1 \pm \zeta)$. In Hartree-Fock, the exponent $\alpha(r_s)$ is a constant, $\alpha = (d+1)/d$ where d is the dimensionality. Hence, for the 2DEG, the Hartree-Fock value of α is 1.5. The coefficients C_1 - C_4 are 1.54039, 0.0305441, 0.296208, and 0.239047 respectively. A comparison of the 2-D and 3-D $\alpha(r_s)$ is given in Fig.1(b).

Our classical mapping of the 2DEG uses the GFMC *spin-polarized* xc-energy fit of Tanatar and Ceperley.⁶ Our unpolarized energies (Table I) are in better agreement with the more recent QMC results of Rapisarda et al.⁷ In Fig. 2 we compare our PDFs with those of GFMC at $r_s = 1, 5$, and 20 given in Tanatar et al, and find satisfactory agreement.

Equation (6) for $p(r_s, \zeta)$ provides a comparison with the results of Varsano et al.,⁸ for the correlation energies at partial polarizations, for $r_s=20$ and 30. Their $-E_c(r_s = 30, \zeta)$ (au.) are, .01216, .01132, .00987, .00776 at $\zeta=.2, .4, .6$ and 0.8, while the results from our fit, eq. (6), are .01219, .01134, .00989, and .00775 respectively. Varsano et al. suggest that the fully spin-polarized phase is more stable, “by an amount close to the statistical uncertainty”. Instead of using the fit, we have evaluated the relevant PDFs accurately at $r_s=30$ and $T = 0$. Varsano et al report a stabilization energy of -1.185×10^{-5} au., while ours is even smaller and not significant.

When r_s is large, the kinetic energy is less impor-

tant; exchange favours spin-polarization and correlation favours the paramagnetic phase. Figure 3 displays the Helmholtz free energy for $r_s = 5, 10, 20$ and 30. Stable spin-polarized phases in the 3DEG have been reported recently.¹⁶ Studies related to the metal-insulator transition in the 2DEG, and numerical studies of 2-D clusters, have increasingly suggested the possibility of ferromagnetic phases of the 2DEG.¹⁷ The 2DEG metal-insulator transition is influenced by the presence of impurities. Our results are for the *pure, uniform* system and there seems to be no stable ferromagnetic phase except near $r_s=30$. For 2DEGs in GaAs, $r_s = 30$ and $T/E_F=0.1$ corresponds to 4.7×10^8 electrons cm^{-2} and 0.019 Kelvin. The more favourable m^* of Si brings the densities close to the $\sim 10^{10}$ electrons cm^{-2} range. Details of the finite- T calculations, the effect of impurities on the spin-polarized phase etc., will be presented elsewhere.

The computer codes used here may be remotely accessed by interested workers by visiting our website.¹⁸

In conclusion, we present a classical calculation of the PDFs, correlation energies etc., of the 2DEG for any spin-polarization and temperature, and find good agreement with quantum-Monte-Carlo results at $T = 0$. The temperature mapping of the quantum effects obtained for 3-D and 2-D systems is found to be universal, to within the accuracy of the available quantum Monte-Carlo exchange-correlations energies. New results for spin-dependent correlation energies and spin polarized distribution functions are given. A comparison of the free energies of the polarized and unpolarized phases at finite- T is presented.

[†] electronic mail address: chandre@cm1.phy.nrc.ca

¹ See, *Electronic properties of two-dimensional systems*, P. Hawrylak, D. J. Lockwood, A. S. Sachrajda, (Editors), Physica E, **6**, 1-892 (2000)

² I. V. Schweigert et al., Phys. Rev. Lett. **82**, 5293 (1999)

³ K. S. Singwi, et al., Phys. Rev. **176** 589 (1968); M. Johnson, J. Phys. C **9**, 3055 (1976)

⁴ D. M. Ceperley, *Recent Progress in Many-body Theories*, Ed. J. B. Zabolitsky (Springer, Berlin 1981), p262

⁵ E. Feenberg, *Theory of Quantum Fluids* (Academic, New York 1969); C. E. Campbell et al., Phys. Rev. B **27**, 7772 (1983); L. J. Lantto, Phys. Rev. B **22**, 1380 (1980)

⁶ B. Tanatar et al., Phys. Rev. B **39**, 5005 (1989)

⁷ F. Rapisarda et al., Aust. J. Phys. **49**, 161 (1996); Y. Kwon et al., Phys. Rev. B **48**, 12037 (1993)

⁸ D. Varsano et al., Europhys. Lett., **53**, 348 (2001)

⁹ M. W. C. Dharma-wardana and F. Perrot, Phys. Rev. Lett. **84**, 959 (2000)

¹⁰ François Perrot and M. W. C. Dharma-wardana, Phys. Rev. B, 15 Dec. (2000)

¹¹ J. M. J. van Leeuwen, J. Grönqveld, J. de Boer, Physica

- 25**, 792 (1959)
- ¹² F. Lado et al Phys. Rev. **A 26**, 2374 (1983) Y. Rosenfeld, Phys. Rev. A **35**, 938 (1987)
- ¹³ M. Minoo et al., Phys. Rev. A **23**, 924 (1981)
- ¹⁴ François Perrot and M. W. C. Dharma-wardana, unpublished.
- ¹⁵ Y. Rosenfeld, Phys.Rev. A **bf42**, 5978 (1990), M. Baus et al., J. Phys. C: Solid State Phys. **19**, L463 (1986)
- ¹⁶ D. P. Young et al., Nature (London) **397**, 412 (1999)
- ¹⁷ S. A. Vitkalov et al., cond-mat/0009454, G. Benenti et al., Phys. Rev. Lett. **86**, 5333 (2001), and A. A. Shashkin et al., *Ibid*, **87**, 86801 (2001)
- ¹⁸ <http://nrcphy1.phy.nrc.ca/ims/qc/chandre/chnc/>


TABLE I. Relevant parameters and a comparison of the QMC correlation energies (au.) of Tanatar and Ceperley (TC), Rapisarda and Senatore (RS), with those from the classical map-HNC (CHNC) for the paramagnetic 2DEG.


r_s	T/T_F	Γ_c	η	$-E_c(\text{TC})$	$-E_c(\text{RS})$	$-E_c(\text{CHNC})$
1	2.000	0.500	0.235	0.10850	- -	0.10838
5	1.849	2.704	0.372	0.04775	0.0489	0.04910
10	1.682	5.946	0.425	0.03034	0.0302	0.03023
20	1.468	13.62	0.469	0.01758	0.0174	0.01739
30	1.331	22.54	0.486	0.01251	0.0124	0.01245


FIG. 1. (a) The quantum temperature T_q for the 2DEG (dashes) and the 3DEG (solid line) as a function of the classical coupling constant $\Gamma = \beta/r_s$. (b) The r_s dependence of the 2DEG (dashes) and 3DEG (solid line) polarization functions at $T = 0$.

FIG. 2. Here the HNC $g(r)$ are compared with GFMC simulations of Tanatar and Ceperley. Solid lines: HNC, boxes: GFMC, Panels (a),(b) unpolarized case, ($\zeta = 0$), $r_s=1$, and $r_s = 20$. Panels (c),(d) $r_s=5$ spin-polarized ($\zeta = 1$) and unpolarized systems.

FIG. 3. The total Helmholtz free energy of the unpolarized ($\zeta = 0$) and polarized ($\zeta = 1$) phases of the 2DEG near $T = 0$ and for $r_s = 5, 10, 20$, and 30, as a function of T/T_F . Solid lines: $\zeta = 0$; dashes with data points (squares): $\zeta = 1$.







Perrot fig.3